



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

$$\begin{aligned}
& \times \sqrt{[a^2c^2 + b^2(x-a)^2]} - a^3(2c^2 - b^2)\sqrt{[b^2 + c^2]} + dc^2(d^2 - 3a^2) \\
& \quad + 3a^2bc^3\log[c(b+d)] - 3a^3bc^2\log a - 3a^2bc^2(x-a)\log[b(x-a)] \\
& + \sqrt{[a^2c^2 + b^2(x-a)^2]} - 3a^3bc^2\log[\sqrt{b^2 + c^2}] - b - 3a^2bc^2(u-a)\log[b(u-a)] \\
& + \sqrt{[a^2c^2 + b^2(u-a)^2]}\} dx du \\
& = \frac{1}{72a^5b^2} \int_0^a \left[24x[2a^2x^2 - 3b^3(x-a)^2]\sqrt{[a^2x^2 + b^2(x-a)^2]} - 16a^3bx^3 \right. \\
& - 16a^3b(x-a)^3 + (x-a)^4(72b^2d - 12a^2d - 12d^3 - 23a^2b) + x^4(12d^3 - 36a^2d \\
& + 23a^2b) + 12a^3x(3b^3 - 2x^2)\sqrt{[b^2 + x^2]} + 12a^3(x-a)[2(x-a)^2 - 3b^2] \\
& \times \sqrt{[b^2 + (x-a)^2]} + 60a^2b(x-a)^4\log[(b+d)(x-a)] \\
& + \frac{24b^4(x-a)^4}{a}\log[(a+d)(x-a)] - 96a^2bx^3(x-a)\log\{b(x-a) \\
& + \sqrt{[a^2x^2 + b^2(x-a)^2]}\} - 48a^3bx^3\log[\sqrt{b^2 + c^2}] - b - \frac{24b^4(x-a)^4}{a} \\
& \times \log\{ax + \sqrt{[a^2x^2 + b^2(x-a)^2]}\} + 48a^3b(x-a)^3\log\{\sqrt{[b^2 + (x-a)^2]} - b\} \\
& + 12a^3b^4\log[x + \sqrt{b^2 + x^2}] - 12a^3b^4\log\{x-a + \sqrt{[b^2 + (x-a)^2]}\} \\
& \left. + 36a^2bx^4\log[x(b+d)] - 48a^3bx^3\log a + 48a^3b(x-a)^3\log a \right] dx \\
& = \frac{1}{16} \left[2d \left(2 - \frac{a^2}{b^2} - \frac{b^2}{a^2} \right) - \frac{a^3 + b^3}{d^2} + \frac{2a^3}{b^2} + \frac{2b^3}{a^2} + \left(\frac{4b^2}{a} - \frac{a^2b^2}{d^3} \right) \log \left(\frac{a+d}{b} \right) \right. \\
& \left. + \left(\frac{4a^2}{b} - \frac{a^2b^2}{d^3} \right) \log \left(\frac{b+d}{a} \right) \right] = \frac{a}{30} [6 + (16 - \sqrt{2})\log(1 + \sqrt{2})], \text{ if } a=b.
\end{aligned}$$

MISCELLANEOUS.

135. Proposed by LON C. WALKER, A. M., Graduate Student, Leland Stanford University, Cal.

Find invariants of the second, third, and sixth degrees in the coefficients of a binary quartic.

Solution by G. B. M. ZERE, A. M., Ph. D., Professor of Chemistry and Physics, The Temple College, Philadelphia, Pa.

Let $ax^4 + 4bx^3y + 6cx^2y^2 + 4dxy^3 + ey^4 = u = 0$, be the binary quartic.

Then $d^4u/dx^4 = 24a$, $d^4u/dx^3dy = 24b$, $d^4u/dx^2dy^2 = 24c$, $d^4u/dxdy^3 = 24d$, $d^4u/dy^4 = 24e$.

Substitute in u , d^4u/dx^4 for y^4 , d^4u/dy^4 for x^4 , $-d^4u/dx^3dy$ for xy^3 , $-d^4u/dxdy^3$ for x^3y , d^4u/dx^2dy^2 for x^2y^2 , and we get $24ae-96bd+144c^2-96bd+24ae=48(ae-4bd+3c^2)$, the invariant of second order.

$$d^2u/dx^2=12(ax^2+2bxy+cy^2).$$

$$d^2u/dy^2=12(cx^2+2dxy+ey^2).$$

$$d^2u/dxdy=12(bx^2+2cxy+dy^2).$$

\therefore The Hessian of this quartic is

$$(ax^2+2bxy+cy^2)(cx^2+2dxy+ey^2)-(bx^2+2cxy+dy^2)^2=0, \text{ or } (ac-b^2)x^4 \\ +2(ad-bc)x^3y+(ae+2bd-3c^2)x^2y^2+2(be-cd)xy^3+(ce-d^2)y^4=0.$$

Write this in the form $Ax^4+4Bx^3y+6Cx^2y^2+4Dxy^3+Ey^4=0$.

Then $24(aE-4bD+6cC-4dB+ Ae)=72(ace+2bcd-ad^2-eb^2-c^3)$, is the invariant of the third order.

$\therefore S=ae-4bd+3c^2$, is the invariant of the second order.

$T=ace+2bcd-ad^2-eb^2-c^3$ is the invariant of the third order.

These are the only two ordinary invariants. The invariant of the sixth order being the discriminant which we determine as below.

If we had used the more general form $Ax^4+By^4+Cz^4$, where $x+y+z=0$, then we would have had $a=A+C$, $e=B+C$, $b=C=e=d$.

$$\therefore S=BC+CA+AB, T=ABC.$$

Equating to zero the two differential equations Ax^3-Cz^3 , and By^3-Cz^3 , we get $Ax^3=By^3=Cz^3$, or $ABCx^3/BC=ABCy^3/AC=ABCz^3/AB$.

$$\therefore x^3 : y^3 : z^3 = BC : AC : AB, \text{ this in } x+y+z=0 \text{ gives } (BC)^{\frac{1}{3}} + (AC)^{\frac{1}{3}} \\ + (AB)^{\frac{1}{3}} = 0.$$

$$\therefore (BC+AC+AB)^3-27A^2B^2C^2=0, \text{ or } S^3-27T^2=0.$$

$$\therefore (ae-4bd+3c^2)^3-27(ace+2bcd-ad^2-eb^2-c^3)^2=\text{the sixth order.}$$

Also solved by G. W. GREENWOOD.

PROBLEMS FOR SOLUTION.

ARITHMETIC.

169. Proposed by F. P. MATZ, Sc. D., Ph. D., Professor of Mathematics and Astronomy in Defiance College, Defiance, Ohio.

At what rate per cent. must a note be discounted at the end of every quarter of a year in order to produce a discount equivalent to 10% interest for the year?